

Mako® Partial Knee

Medial PKA patient indications



Who and why?

Patient candidacy continues to be one of the most common concerns for partial knee arthroplasty. Below, see how surgeons evaluated various patient histories to identify candidates for a medial Mako Partial Knee.



David Howe, M.D.


Patient history

- 69 year old male
- Active; plays golf and pickleball; cycles daily
- Right knee pain for 2+ years
- Medial bone on bone osteoarthritis
- Isolated unicompartmental disease

Preop X-rays



Postop X-rays



Michael Morwood, M.D.


Patient history

- 67 year old female
- Active
- Right knee pain
- Significant medial joint space narrowing
- Retained lateral joint space and appropriate patella tracking

Preop X-rays



Postop X-rays






Akhil Sastry, M.D.


Patient history

- 83 year old female with a BMI of 26
- Long standing pain along medial aspect of knee for 2 years
- Has tried PT and injections with no improvement
- Range of motion 0-120 degrees
- Negative Lachman test

Preop X-rays



Postop X-rays



Ismar Dizdarevic, M.D.


Patient history

- Active 68 year old male
- Medial left knee pain for 1 year
- Attempted meniscus debridement 10 months prior with no improvement
- Correctable varus
- Negative patellar grind 0-135 degrees of motion

Preop X-rays



Postop X-rays



The faculty are paid consultants of Stryker Orthopaedics. The opinions expressed by the faculty are those of the faculty and not necessarily those of Stryker.

A surgeon must always rely on his or her own professional clinical judgment when deciding whether to use a particular product when treating a particular patient. Stryker does not dispense medical advice and recommends that surgeons be trained in the use of any particular product before using it in surgery. The information presented is intended to demonstrate the breadth of Stryker's product offerings. A surgeon must always refer to the package insert, product label and/or instructions for use before using any of Stryker's products. Products may not be available in all markets because product availability is subject to the regulatory and/or medical practices in individual markets. Please contact your sales representative if you have questions about the availability of products in your area.

Stryker Corporation or its divisions or other corporate affiliated entities own, use or have applied for the following trademarks or service marks: Mako, Stryker. All other trademarks are trademarks of their respective owners or holders.